60th Medical Group (AMC), Travis AFB, CA

INSTITUTIONAL ANIMAL CARE AND USE COMMITTEE (IACUC)

FINAL REPORT SUMMARY

(Please type all information. Use additional pages if necessary.)

PROTOCOL #: FDG20130029A			DATE: 2 December 2013					
	OCOL TITLE: Compa injuries	arative testing of hemostatic of	dressings in a large animal m	odel (Sus scrofa) with severe				
PRINCI	IPAL INVESTIGATO	R (PI) / TRAINING COORDI	INATOR (TC): Capt Hilary G	allogly				
DEPAR	RTMENT: Clinical In	vestigations Facility	PHONE #: 423-7400					
INITIAL	. APPROVAL DATE	: 24 April 2013	LAST TRIENNIAL REVISION DATE: N/A					
FUNDI	NG SOURCE: Air	Force Surgeon General's Off	fice					
1.	RECORD OF ANIM	IAL USAGE:						
An	imal Species:	Total # Approved	# Used this FY	Total # Used to Date				
Sus scr	rofa	36	18	18				
Note. Many fewer animals than approved were used because one of the original treatment groups (Lypressin-soaked gauze) could not be done due to the drug not being available. The remaining 3 treatment groups had only 6 animals per group instead of 9 because it became apparent that there was no difference between treatments and early stopping rules were applied.								
2.	Training: Live Training: non-l	Live Animal Heavival (chronic) Precon-Survival (acute) Util	dical Readiness alth Promotion evention	Prolonged Restraint Multiple Survival Surgery Behavioral Study Adjuvant Use				
3.	PROTOCOL PAIN CATEGORY (USDA): (Check applicable) CX_D E							
4.	PROTOCOL STATUS:							
	*Request Protocol Closure:							
	Inactive, protocol never initiated							
	Inactive, protocol initiated but has not/will not be completed							
	X Completed, all approved procedures/animal uses have been completed							
5.	FUNDING STATUS	E: Funding allocated:	\$20,160.00 Fur	nds remaining: \$ 0.00				
6.	PROTOCOL PERS	ONNEL CHANGES:						
	nere been any perso nal review?	nnel/staffing changes (PI/CI/A _x_ Yes No		st IACUC approval of protocol,				

Report Documentation Page

Form Approved OMB No. 0704-018

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 19 DEC 2013	2. REPORT TYPE Final	3. DATES COVERED 24 Apr 2013 - 19 Dec 2013	
4. TITLE AND SUBTITLE	5a. CONTRACT NUMBER		
FDG20130029A "Comparative testing animal model (Sus Scorofa) with seve	5b. GRANT NUMBER		
ammai model (Sus Scorola) with seve	5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) Capt. Hilary Gallogly	5d. PROJECT NUMBER FDG20130029A		
	5e. TASK NUMBER		
	5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND A Clinical Investigation Facility David Circle Travis AFB, CA 94535	8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S)	10. SPONSOR/MONITOR'S ACRONYM(S)		
Clinical Investigation Facility David Circle Travis AFB, CA 94535	11. SPONSOR/MONITOR'S REPORT NUMBER(S)		

12. DISTRIBUTION/AVAILABILITY STATEMENT

Approved for public release, distribution unlimited

13. SUPPLEMENTARY NOTES

14. ABSTRACT

OBJECTIVES: We compared the hemostatic and hemodynamic effects of dilute concentrations of epinephrine soaked gauze and with epinephrine soaked Combat Gauze (CG) in swine with grade IV liver injuries. METHODS: Anesthetized swine were instrumented, splenectomized, and had a grade IV liver injury created. After 30 seconds of free bleeding, damage control liver packing was performed with laparotomy pads soaked in different concentrations of epinephrine (1 mg/L or 2mg/L normal saline) or CG pads soaked in 3mg epinephrine/L normal saline. Hemodynamic and laboratory data were recorded, and blood loss was measured for two hours. Post-mortem histopathology was performed on the liver injury sites. RESULTS: There were no pre-injury differences between groups, and all animals survived the entire two hours. Animals treated with 1 mg/L epi, 2 mg/L epi, and 3 mg/L epi-CG dressings had similar amounts of blood loss (18.7, 22.3, and 21.8 mL/kg respectively, p = 0.85). There were no significant differences between groups in laboratory measurements or physiology measurements. Histopathology revealed no adverse cellular effects from any treatment. CONCLUSION: There were no significant or practical differences in blood loss from animals treated with 1 mg/L and 2 mg/L epinephrine. In a previous experiment, we found that 3 mg/L epinephrine soaked gauze resulted in much less blood loss (16.5 mL/kg) and those animals had similar heart rates and mean arterial pressures as those we observed in this study. Using the more concentrated epinephrine solution with Combat Gauze did not offer any advantage over plain gauze with epinephrine. Therefore, we recommend using a more concentrated epinephrine solution (3 mg epinephrine per liter sterile saline) as a hemostatic dressing in severe liver injuries.

15. SUBJECT TERMS

US Air Force, Medical Service, Medical Research, Graduate Medical Education

16. SECURITY CLASSIFIC	17. LIMITATION OF	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON		
a. REPORT unclassified	ь. abstract unclassified	c. THIS PAGE unclassified	UU	3	RESPONSIBLE PERSON

Standard Form 298 (Rev. 8-98) Prescribed by ANSI Std Z39-18 If yes, complete the following sections (Additions/Deletions). For additions, indicate whether or not the IACUC has approved this addition.

ADDITIONS: (Include Name, Protocol function - PI/CI/AI/TC/Instructor, IACUC approval - Yes/No)

Capt Paul Vu, Al-Surgeon, approved by the IACUC on 9 September 2013.

<u>DELETIONS</u>: (Include Name, Protocol function - PI/CI/AI/TC/Instructor, Effective date of deletion)

PROBLEMS / ADVERSE EVENTS: Identify any problems or adverse events that have affected study progress. Itemize adverse events that have led to unanticipated animal illness, distress, injury, or death; and indicate whether or not these events were reported to the IACUC.

None.

8. REDUCTION, REFINEMENT, OR REPLACEMENT OF ANIMAL USE:

REPLACEMENT (ALTERNATIVES): Since the last IACUC approval, have alternatives to animal use become available that could be substituted in this protocol without adversely affecting study or training objectives?

No.

REFINEMENT: Since the last IACUC approval, have any study refinements been implemented to reduce the degree of pain or distress experienced by study animals, or have animals of lower phylogenetic status or sentience been identified as potential study/training models in this protocol?

No.

REDUCTION: Since the last IACUC approval, have any methods been identified to reduce the number of live animals used in this protocol?

No.

9. PUBLICATIONS / PRESENTATIONS: (List any scientific publications and/or presentations that have resulted from this protocol. Include pending/scheduled publications or presentations).

Submitted for presentation at the Society of Air Force Clinical Surgeons annual meeting.

- 10. Were the protocol objectives met, and how will the outcome or training benefit the DoD/USAF?
- Yes. The protocol outcomes will help guide therapeutic choices for Air Force surgeons operating on liver injuries.
- **11. PROTOCOL OUTCOME SUMMARY:** (Please provide, in "ABSTRACT" format, a summary of the protocol objectives, materials and methods, results include tables/figures, and conclusions/applications.)

OBJECTIVES: We compared the hemostatic and hemodynamic effects of dilute concentrations of epinephrine soaked gauze and with epinephrine soaked Combat Gauze™ (CG) in swine with grade IV liver injuries.

METHODS: Anesthetized swine were instrumented, splenectomized, and had a grade IV liver injury created. After 30 seconds of free bleeding, damage control liver packing was performed with laparotomy pads soaked in different concentrations of epinephrine (1 mg/L or 2mg/L normal saline) or CG pads soaked in 3mg epinephrine/L normal saline. Hemodynamic and laboratory data were recorded, and blood loss was measured for two hours. Postmortem histopathology was performed on the liver injury sites.

RESULTS: There were no pre-injury differences between groups, and all animals survived the entire two hours. Animals treated with 1 mg/L epi, 2 mg/L epi, and 3 mg/L epi-CG dressings had similar amounts of blood loss (18.7, 22.3, and 21.8 mL/kg respectively, p = 0.85). There were no significant differences between groups in laboratory measurements or physiology measurements. Histopathology revealed no adverse cellular effects from any treatment.

CONCLUSION: There were no significant or practical differences in blood loss from animals treated with 1 mg/L and 2 mg/L epinephrine. In a previous experiment, we found that 3 mg/L epinephrine soaked gauze resulted in

much less blood loss (16.5 mL/kg) and those animals had similar heart rates and mean arterial pressures as those we observed in this study. Using the more concentrated epinephrine solution with Combat Gauze™ did not offer any advantage over plain gauze with epinephrine. Therefore, we recommend using a more concentrated epinephrine solution (3 mg epinephrine per liter sterile saline) as a hemostatic dressing in severe liver injuries.

(Principal Investigator Signature

20 February 2014 (Date)